

Endogenous FVIII activity and procedure-related FVIII use and bleeding: Post hoc analysis of GENE8-1

*Doris V Quon¹, Jiaan-Der Wang², Michael Wang³, Dominic Pepperell⁴, Young Shil Park⁵, Gili Kenet⁶, Johnny Mahlangu⁷, Teh-Liane Khoo⁸, **Simon Harris⁹**, Tara M Robinson⁹, Konstantia-Maria Chavele⁹, Steven W Pipe¹⁰*


¹Orthopaedic Hemophilia Treatment Center, Los Angeles, CA, USA; ²Center for Rare Disease and Hemophilia, Taichung Veterans General Hospital, Taichung, Taiwan; ³Hemophilia and Thrombosis Center, University of Colorado Anschutz Medical Campus, Aurora, CO, USA; ⁴Department of Haematology, Fiona Stanley Hospital, Murdoch, WA, Australia; ⁵Kyung Hee University Hospital at Gangdong, Seoul, South Korea; ⁶The National Hemophilia Center, and Amalia Biron Research Institute of Thrombosis and Hemostasis, Sheba Medical Center, Tel Hashomer, Tel Aviv University, Tel Aviv, Israel; ⁷Hemophilia Comprehensive Care Center, Charlotte Maxeke Johannesburg Academic Hospital, University of the Witwatersrand and NHLS, Johannesburg, South Africa; ⁸Haematology Department, Haemophilia Treatment Centre, The Royal Prince Alfred Hospital, Sydney, NSW, Australia; ⁹BioMarin Pharmaceutical Inc., Novato, CA, USA; ¹⁰Departments of Pediatrics and Pathology, University of Michigan, Ann Arbor, MI, USA

Disclosures for Simon Harris

Research support/principal investigator	No relevant conflicts of interest to declare
Employee	Employee and stockholder of BioMarin Pharmaceutical Inc.
Consultant	No relevant conflicts of interest to declare
Major stockholder	No relevant conflicts of interest to declare
Speakers bureau	No relevant conflicts of interest to declare
Honoraria	No relevant conflicts of interest to declare
Scientific advisory board	No relevant conflicts of interest to declare

Evaluating endogenous FVIII activity and procedure-related FVIII use following 2 years of treatment with valoctocogene roxaparvovec

- Valoctocogene roxaparvovec (AAV5-hFVIII-SQ) transfers a FVIII coding sequence to hepatocytes using a recombinant AAV5 vector, enabling endogenous FVIII production in people with severe haemophilia A¹⁻⁴
- In GENE8-1, an open-label, phase 3 trial, participants achieved FVIII activity providing improved protection from bleeding compared with prophylaxis for 52 and 104 weeks^{4,5}
- Here, we present findings from a post hoc analysis of procedures in GENE8-1 after 2 years of follow-up, with a focus on FVIII use, bleeding outcomes, and rationale for PI decisions




AAV, adeno-associated virus serotype 5; FVIII, factor VIII; hFVIII-SQ, human FVIII, SQ variant; PI, principal investigator.

1. Rangarajan S, et al. *N Engl J Med.* 2017;377(26):2519-30; 2. Pasi KJ, et al. *N Engl J Med.* 2020;382(1):2940; 3. Pasi KJ, et al. *Haemophilia.* 2021;27(6):947-56; 4. Ozelo M, et al. *N Engl J Med.* 2022;386(11):1013-25; 5. Mahlangu J, et al. 15th Annual Congress of EAHAD. February 2–4, 2022.

Phase 3 GENE8-1 study design

Eligible participants

- Participants directly enrolled or were rolled over from the non-interventional study BMN 270-902
- Adult men with severe haemophilia A (FVIII ≤ 1 IU/dL)
- Previously receiving FVIII prophylaxis
- No history of FVIII inhibitors or anti-AAV5 antibodies



^aFVIII activity measured by the chromogenic substrate assay.
AAV5, adeno-associated virus serotype 5; FVIII, factor VIII; W, week.

Baseline demographics and characteristics of ITT population

Baseline characteristics ¹	ITT n=134
Age, years, mean ± SD	31.7 ± 10.3
Race, n (%)	
White	96 (71.6)
Asian	19 (14.2)
Black or African American	15 (11.2)
Hawaiian or Pacific Islander	1 (0.7)
Not provided	3 (2.2)
Hispanic or Latino ethnicity, n (%)	7 (5.2)
BMI, kg/m², mean ± SD	25.3 ± 4.6
Medical history, n (%)	
Hepatitis B	20 (14.9)
Hepatitis C	41 (30.6)
HIV	2 (1.5)
Number of problem joints,^a n (%)	
0	97 (72.4)
1	17 (12.7)
2	9 (6.7)
3	8 (6.0)
>3	3 (2.2)


^aProblem joints were those with chronic joint pain, chronic synovitis, haemophilic arthropathy, limited motion, or recurrent bleeding.

BMI, body mass index; HIV, human immunodeficiency virus; ITT, intent to treat; SD, standard deviation.

1. Ozelo M, et al. *N Engl J Med.* 2022;386(11):1013-25.

Procedures performed during GENE8-1

- In total, 77 participants underwent 260 total procedures
- Types of procedures¹
 - Non-invasive: eg, tattoo, dental cleaning
 - Invasive:
 - Major: eg, joint debridement, arthrodesis
 - Minor: eg, dental extraction, biopsies
- PIs were sent questionnaires asking what factors influenced their decision to perform procedures without FVIII infusions





Perioperative management for participants in GENE8-1

- Major invasive procedures were performed with FVIII treatment regardless of participant FVIII level
- Compared with minor procedures, major procedures were associated with a higher:
 - FVIII dose
 - Number of FVIII infusions
 - Days of post-operative FVIII treatment


	Minor invasive procedures	Major invasive procedures
Mean FVIII dose, IU/kg (min/max)	67.2 (14–324)	255.4 (103–538)
Number of FVIII infusions, n (min/max)	2.2 (1–13)	8.8 (3–21)
Days of post-operative FVIII, n (min/max)	1.8 (1–7)	6.9 (2–14)

Minor invasive procedures performed without FVIII treatment were associated with higher participant FVIII activity per CSA




Relatively few invasive procedure–related bleeds occurred




- There were 18 procedure-related bleeding episodes in 14 participants
- Bleeding episodes were self-reported by participants
 - 13 required FVIII treatment
 - Mean FVIII activity: 10.4 IU/dL
 - Median FVIII activity: 6.9 IU/dL
 - 5 did not require FVIII treatment
 - Mean FVIII activity: 60.4 IU/dL
 - Median FVIII activity: 52.1 IU/dL




Effective haemostatic control was possible for participants of the GENE8-1 trial




procedures were performed without FVIII treatment



(32.8%) participants had pre-operative FVIII activity below 20 IU/dL



PIs responded to a questionnaire asking why they didn't treat participants with FVIII



PIs said they deemed the participant's endogenous FVIII expression sufficient for the procedure

Conclusions

- In the GENE8-1 clinical trial of valoctocogene roxaparvovec for severe haemophilia A, 67 out of 111 invasive procedures were performed without FVIII treatment
 - Most minor invasive procedures (62/67) were safely performed without bleeding events
 - For minor invasive procedures, FVIII treatment was associated with lower participant pre-operative endogenous FVIII activity
- PI questionnaire responses reflect the nature of personalized medicine with valoctocogene roxaparvovec
 - Most PIs and participants had case-by-case discussions about the use of FVIII treatment for procedures, given the participant's endogenous FVIII activity achieved with gene therapy and the type of procedure
- The data from this study, and the opinions reflected in the PI questionnaire, suggest that valoctocogene roxaparvovec provides effective haemostatic control for up to 2 years even with FVIII activity below WFH guidelines



Acknowledgements

**Thank you to all the trial participants, their families,
study site personnel, and investigators**

- Funding for this study was provided by BioMarin Pharmaceutical Inc.
- Medical writing and editorial support were funded by BioMarin Pharmaceutical Inc.