Systematic literature review of prevalence and severity of somatic comorbidities in adults with PKU

Harding C¹, Whitehall KB², Rose S², Clague GE², Ahring KK³, Bilder DA⁴, Hermida Á⁵, Inwood A⁶, Longo N⁷, Maillot F⁸, Muntau AC⁹, Pessoa ASL¹⁰, Rocha JC¹¹, Rohr F¹², Sivri S¹³, Said J¹⁴, Oshinbolu S¹⁵, Sibbring GC¹⁴

¹Oregon Health & Science University, Portland, OR, USA; ²BioMarin Pharmaceutical Inc., Novato, CA, USA; ³Center for PKU, Copenhagen University Hospital, Copenhagen, Denmark; ⁴Huntsman Mental Health Institute, University of Utah, Salt Lake City, UT, USA; 5University of Santiago de Compostela, Santiago de Co Utah School of Medicine, Salt Lake City, UT, USA; 8CHRU and the University of Tours, France; 9University Medical Center Hamburg-Eppendorf, Hamburg, Germany; 10Ceará State University and Hospital Infantil Albert Sabin, Fortaleza, CE, Brazil; ¹¹NOVA Medical School, NOVA University of Lisbon, and Reference Centre Inherited Metabolic Diseases, Centro Hospitalar Universitário Lisboa Central, Lisbon, Portugal; ¹²Met Ed Consultants, Boulder, CO, USA; ¹³Hacettepe University, Ankara, Turkey; ¹⁴Prime Global, Knutsford, UK; ¹⁵Prime Global, London, UK


COPD = chronic obstructive pulmonary disease.

Introduction

- Phenylketonuria (PKU) is an inborn error of phenylalanine (Phe) metabolism that, if untreated, causes Phe accumulation in the blood and brain leading to neurophysiological alterations and poor developmental outcomes¹
- Lifelong management of PKU centers on medical nutrition therapy with a Phe-restricted diet to achieve (and maintain) appropriate blood Phe levels^{2,3}; sapropterin dihydrochloride is indicated for responsive individuals^{4,5} and pegvaliase is an option for adults with uncontrolled blood Phe levels despite intervention^{6,7}
- Many adults are unable to achieve blood Phe levels within guideline-recommended ranges, either due to waning adherence to a Phe-restricted diet or an inadequate response to treatment, including due to disease severity^{8,9}; uncontrolled blood Phe levels lead to adverse neurocognitive and neuropsychiatric outcomes¹⁰⁻¹²
- Comorbidities across organ systems have been reported in adults with PKU, with claimsbased studies finding a higher prevalence of somatic comorbidities compared with a general population^{13,14}
- A systematic literature review (SLR) was conducted to evaluate the prevalence and severity of non-neurocognitive and non-neuropsychiatric comorbidities in adults with PKU by intervention, disease severity, and adherence, to provide insight into specific treatment situations

Methods

- The SLR is registered with the Research Registry (reviewregistry1476)
- Eligibility criteria were established using the Population, Intervention, Comparator, Outcome, Study design (PICOS) framework (Figure 1)



Inclusion: No therapeutic intervention (not receiving medical nutritional therapy/ Phe-restricted diet, and/or sapropterin dihydrochloride or pegvaliase, including healthy controls or a reference population) Protein substitutes and/or low protein foods

Inclusion: The prevalence or severity of different somatic comorbidities in patients with PKU on medical nutritional therapy/Phe-restricted diet, and/or sapropterin dihydrochloride or pegvaliase compared with: healthy controls the general population (including standard reference values) patients with PKU not receiving any form of therapeutic intervention patients with PKU who did not adhere to treatment patients with PKU who interrupted/discontinued treatment The prevalence or severity of different somatic comorbidities in patients across the PKU disease spectrum on medical nutritional therapy/Phe-restricted diet, and/or sapropterin dihydrochloride or pegvaliase

Inclusion: Randomized controlled trials; single-arm clinical trials; cohort studies (prospective and retrospective); cross-sectional studies and surveys **Exclusion:** Systematic reviews and meta-analyses; narrative (non-systematic) review articles; animal or in vitro studies; letters, editorials, and commentaries; guidelines and best practice; congress abstracts; non-peer-reviewed articles Inclusion: Language of publication: English; date of publication: up to February 1, 2022; countries: all

^aStudies that included children but did not stratify outcomes by patient age were included.

- Literature was retrieved via the PubMed interface and included publications from MEDLINE from earliest coverage (1946) to February 1, 2022, using pre-defined free text and Medical Subject Headings (MeSH) search terms based on the PICOS framework
- A two-stage screening process identified records eligible for data extraction

Exclusion: Language of publication: non-English


- Records were screened by abstract, and those considered potentially eligible were screened by full text to confirm eligibility (concordance of eligibility decisions was assessed by independent review of 10% of records selected at random)
- Data extraction was conducted by one reviewer into a pre-designed spreadsheet (Microsoft) Excel®) and was checked for accuracy by an independent reviewer
- Studies were grouped by PKU populations identified and by comorbidity types Vote counting was used for data synthesis, allowing direction of effect to be determined
- Vote counting followed the methods described in the Cochrane handbook¹⁵ and was reported according to the Synthesis Without Meta-analysis (SWiM) guidelines¹⁶
- A standardized binary metric was created by allocating votes to the groups compared in studies according to the direction of a higher comorbidity burden, regardless of statistical significance of differences between the groups
- The number of votes allocated to each group was then compared to determine the direction of effect

Results

Study selection and characteristics

- In total, 1,128 unique records were screened; 459 records were identified as potentially eligible for inclusion and 73 studies spanning 10,640 patients were confirmed as eligible for inclusion (Figure 2)
- Most were of observational design and most were conducted in European countries (n=53) and/or in North America (n=19)

Figure 2. PRISMA diagram showing article selection process Identification of studies via databases



PRISMA = Preferred Reporting Items for Systematic Reviews and Meta-Analyses. Articles were excluded on a hierarchical basis, in the order that questions were asked (i.e., if the answer to the first question was no, this was given as the main reason for exclusion, but articles may have met or not met other criteria). alncludes studies that did not present outcomes in a meaningful way which answered one or more of the pre-specified research questions. Includes open interventional trials, pooled analyses, and cost analyses.

- More than 15 different comorbidity types were reported across the 73 studies: white matter abnormalities and bone-related abnormalities were the most commonly reported
- Studies were grouped according to PKU populations (Table 1) and then according to comorbidity type

Table 1. Grouping of studies by PKU population

| Study groupings | | Number of | Number of | Vote counting |
|---|--|----------------------------------|-----------------------|------------------|
| Intervention population | Comparator population | studies | patients ^a | conducted |
| Patients with PKU on a Phe-restricted diet with or without pharmacologic therapy | Healthy controls or reference values | 58 ^{13,14,17-72} | 9,938 | Yes ^b |
| Patients with PKU adherent to a Phe-restricted diet | Patients with PKU non-adherent to a Phe-restricted diet | 88,44,65,73-77 | 417 | Yes ^c |
| Patients with PKU on a specific Phe-restricted diet | Patients with PKU on a different Phe-restricted diet | 4 ⁷⁸⁻⁸¹ | 96 | Nod |
| Patients with a more severe PKU on a Phe-restricted diete | Patients with HPA or a less severe PKU on a Phe-restricted diet ^e | 382-84 | 123 | Nod |
| Patients with PKU treated with pegvaliase or sapropterin dihydrochloride ^f | Patients with PKU treated with placebo | 285,86 | 166 | Nod |

HPA = hyperphenylalaninemia ^aTwo studies^{44,65} are counted twice as both compare patients with PKU on a Phe-restricted diet with or without pharmacologic therapy versus healthy controls or reference values, and patients with PKU who adhered to a Phe-restricted diet versus non-adherent. Vote counting was conducted on 47 studies: nine of the 58 studies did not report results for the healthy controls group and were excluded; another two studies were also excluded as it was not possible to confirm treatment with a Phe-restricted diet in the full study population. Vote counting was conducted on seven studies: one study was excluded due to no clear correlation between the severity of comorbidity and dietary history of the patients. Fewer than five studies resulting in insufficient data, vote counting could not be conducted. eIncludes studies of patients with classical PKU versus patients with mild/moderate PKU and patients with PKU versus patients with HPA. One study included five participants who followed a Phe-restricted diet (study did not specify whether these patients were in the pegvaliase or placebo group). Another study⁸⁶ did not indicate patients were on a Phe-restricted diet.

Data synthesis

- In the vote-counting analysis shown in Figure 3, a higher burden of ≥1 comorbidity (or outcome measure) in patients with PKU versus healthy controls or reference values was indicated in all 47 studies, and a higher burden of ≥1 comorbidity (or outcome measure) in healthy controls or reference values versus patients with PKU was indicated in seven studies
- Bone-related abnormalities and white matter abnormalities were the most commonly reported comorbidities with a higher burden of ≥1 comorbidity (or outcome measure) in patients with PKU on a Phe-restricted diet with or without pharmacologic therapy versus healthy controls or reference values

Figure 3. Burden of comorbidities in patients with PKU on a Phe-restricted diet with or without pharmacologic therapy versus healthy controls or reference values (assessed by vote counting, n=47 studies)






Figure indicates the number of studies with a higher burden of ≥1 comorbidity or outcome measure (some studies reported more than one comorbidity/outcome measure or had a differing direction of effect between comorbidities/outcome measures). Vote counting was conducted regardless of statistical significance. In most studies, all patients were on a Phe-restricted diet, except: one study with a mixture of patients on and not on a Phe-restricted diet23; one study in which some patients received sapropterin, some patients were on a Phe-restricted diet, and, for some patients, it was not clear whether they were on a Phe-restricted diet or not¹⁴; and one study in which some patients were treated with sapropterin in addition to dietary treatment³⁵. Bone-related abnormalities: Higher burden in PKU (n=19)^{14,17,22,23,26,28,29,33,40,42-45,50,56,57,63,70,71}; higher burden in controls (n=4)^{17,23,29,45}. White matter abnormalities: Higher burden in PKU (n=13)30,31,34,39,41,49,52-55,64,66,69. Cardiovascular outcomes: Higher burden in PKU (n=4)14,18,35,36; higher burden in controls (n=1)36. Nutritional outcomes: Higher burden in PKU (n=4)37,38,67,68. Overweight/obesity: Higher burden in PKU (n=3)14,58,60; higher burden in controls (n=2)58,60. Diabetes: Higher burden in PKU (n=2)²⁷. Tremors: Higher burden in PKU (n=2)^{47,53}. Hypertension: Higher burden in PKU (n=1)¹⁴. Dermatologic disorders: Higher burden in PKU (n=1)¹⁴. Gastrointestinal disorders: Higher burden in PKU (n=1)14. COPD/asthma: Higher burden in PKU (n=1)14. Other: Higher burden in PKU (n=6)14,20,23-25,65; higher burden in controls $(n=2)^{23,60}$.

- A range of outcome measures for bone-related abnormalities and white matter abnormalities was reported across studies (Figure 4)
- The most commonly reported (in ≥4 studies) outcome measures for bone-related abnormalities were Z-scores, markers for bone resorption and bone formation, and prevalence of osteopenia/ osteoporosis
- The most commonly reported (in ≥4 studies) outcome measures for white matter abnormalities were magnetic resonance imaging grading, apparent diffusion coefficient, and fractional anisotropy
- In the vote-counting analysis comparing the burden of comorbidities in patients with PKU adhering to a Phe-restricted diet versus those non-adherent, four studies indicated a higher burden of ≥1 comorbidity (or outcome measure) in patients who adhered versus those non-adherent, and six studies indicated a higher burden of ≥1 comorbidity (or outcome measure) in patients who did not adhere versus those adherent
- A range of comorbidities was reported in the other PKU population comparisons:
- White matter abnormalities were reported across PKU severity types in a single study of patients with classical PKU versus mild disease (including a mixture of patients who were early and late-treated, patients off diet, and patients on a Phe-restricted diet); no significant difference was found in terms of the severity of white matter involvement between the disease severity groups $(p=0.36)^{82}$
- Bone-related abnormalities were reported in another study of patients who were on a Phe-restricted diet with classical PKU versus mild/moderate disease; the prevalence of osteopenia and osteoporosis was reported to be similar between the disease severity groups (no statistical comparison reported)83
- Nutritional outcomes were measured in patients with PKU and patients with HPA; differences in the concentrations of serum pre-albumin, zinc, and iron between groups were not statistically significant, but there was a statistically significant lower concentration of selenium in adult patients with PKU compared with adult patients with HPA (p=0.006)84
- Headache was reported as an adverse event in studies of pegvaliase or sapropterin dihydrochloride^{85,86} - One study of patients with PKU reported more headaches in the placebo group than in the pegvaliase group (no statistical comparison reported)85
 - Another study reported headaches in patients with PKU or HPA receiving sapropterin dihydrochloride; headaches (in one patient), which were considered to be probably related to sapropterin and headaches (in eight patients) and migraines (in four patients), which were considered to be possibly related to sapropterin⁸⁶
- Three studies evaluating different Phe-restricted diets reported on prevalence of overweight/ obesity⁷⁸⁻⁸⁰. Two studies reported no significant differences between diets (p=1.000⁷⁹, p=0.367⁸⁰); one study reported body mass index remained unchanged between diets⁷⁸

Figure 4. Measures used to report A) bone-related abnormalities and B) white matter abnormalities in patients with PKU on a Phe-restricted diet with or without pharmacologic therapy versus healthy controls or reference values



BAP = bone alkaline phosphatase; BMD = bone mineral density; ICTP = pyridinoline cross-linked telopeptide domain of type I collagen; OC = osteoclastogenesis; PR = prevalence ratio. Underlined studies showed a statistically significant difference between groups. All 20 studies indicated a higher prevalence and/or severity in the PKU group compared with healthy controls or reference values; with 17 reporting a statistically significant difference 13,14,17,22,23,28,33,40,42-45,50,56,57,63,70, one that did not find a statistically significant difference²⁶, and two that did not test for statistical significance between PKU group and controls^{29,71}. Further details on this figure can be found in the forthcoming manuscript.



ADC = Apparent diffusion coefficient; IW = intermediate weighted; NMR = nuclear magnetic resonance. Out of 19 studies, seven studies did not report outcome values for healthy controls despite a clear indication of a control group, and therefore the results are not comparative^{21,32,54,61,62,72}. From the 12 studies that did report outcome values for both healthy controls/reference values and patients with PKU^{30,31,34,39,41,46,49,52,53,55,66,69}. all 12 indicated a higher prevalence and/or severity in the PKU group compared with healthy controls or reference values; with eight studies reporting statistical significance^{30,31,34,39,49,52,66,69} and four studies that did not test for statistical significance^{41,53,55,64}. Underlined studies showed a statistically significant difference between groups. Further details on this figure can be found in the forthcoming manuscript

Conclusions

- This SLR provides evidence for an unmet need in the current treatment landscape, highlighting the higher somatic, non-neurocognitive and non-neuropsychiatric comorbidity burden on patients with PKU versus a non-PKU population. To potentially avoid the clinical and economic implications of managing comorbidities, there is a need for improved access to therapeutic interventions to maintain blood Phe levels within recommended ranges over the long term
- No conclusions could be drawn from vote counting of the diet adherent versus non-adherent populations, as numbers of studies in each population indicating a higher burden were similar
- The scarcity of data from studies using similar designs and patient populations, as well as consistency of outcome measures for many comorbidities, restricted the synthesis methods that could be used to evaluate the somatic comorbidity burden
 - Although vote counting is considered an acceptable alternative when meta-analysis is not feasible¹⁵, not all studies could be included in the analysis due to lack of comparative data; the analysis does not account for differences in the relative sizes of the studies or methodological aspects, and provides no information on the magnitude of effect
- More robust studies reporting consistent outcome measures and evaluating the relationship between effective metabolic control and comorbidity burden are needed

References

1. Williams RA et al. Clin Biochem Rev. 2008;29:31-41. 2. Vockley J et al. Genet Med. 2014;16:188-200. 3. van Wegberg AMJ et al. Orphanet J Rare Dis. 2017;12:162. 4. BioMarin Pharmaceutical Inc. KUVAN Summary of Product Characteristics. 2020. Available at. https://www.ema.europa.eu/en/documents/product-information/kuvan-epar-product-information_en.pdf. Accessed January 19, 2023. 5. BioMarin Pharmaceutical Inc. KUVAN Prescribing information. 2021. Available at: http://kuvan.com/wp-content/file/KUVAN_ Prescribing_Information1.pdf. Accessed January 19, 2023. 6. BioMarin Pharmaceutical Inc. PALYNZIQ Summary of Product Characteristics. 2022. Available at: https://www.ema.europa.eu/en/documents/product-information/palynziq-epar-product-information_ en.pdf. Accessed January 19, 2023. 7. BioMarin Pharmaceutical Inc. PALYNZIQ Prescribing information. 2020. Available at: https://www.palynziq.com/prescribinginformation.pdf. Accessed January 19, 2023. 8. Green B et al. Nutrients. 2019;11. 9. Kanufre V et al. Nutrients. 2021;13:3118. 10. Pilotto A et al. Neurology. 2021;96:e399-e411.11. Bilder DA et al. Neuropsychol. 2016;41:245-260. 12. Romani C et al. Neurosci Biobehav Rev. 2022;143:104925. 13. Burton BK et al. Mol Genet Metab. 2018;125:228-234. 14. Trefz KF et al. Orphanet J Rare Dis. 2019;14:181. 15. McKenzie JE et al. Cochrane Handbook for Systematic Reviews of Interventions. Chapter 12: Synthesizing and presenting findings using other methods. 2022. Available at: https://training.cochrane.org/ handbook/current/chapter-12. Accessed January 19, 2023. 16. Campbell M et al. BMJ. 2020;368:16890. 17. Adamczyk P et al. J Bone Miner Metab. 2011;29:236-244. 18. Azabdaftari A et al. Orphanet J Rare Dis. 2019;14:213. 19. Bilder DA et al. Mol Genet Metab. 2017;121:1-8. 20. Bodner KE et al. Mol Genet Metab. 2012;107:302-307. 21. Burlina AB et al. J Inherit Metab Dis. 2000;23:313-316. 22. Carson DJ et al. Pediatr Radiol. 1990;20:598-599. 23. Choukair D et al. J Inherit Metab Dis. 2017;40:219-226. 24. Christ SE et al. Mol Genet Metab. 2010;99 Suppl 1:S33-40. 25. Christ SE et al. Mol Genet Metab. 2016;118:3-8. 26. Coakley KE et al. J Inherit Metab Dis. 2016;39:363-372. 27. Couce ML et al. Orphanet J Rare Dis. 2018;13:103. 28. de Groot MJ et al. Mol Genet Metab 2012;105:566-570. **29**. Demirdas S et al. Ann Nutr Metab. 2017;70:111-121. **30**. Dezortová M et al. Acta Radiol. 2001;42:459-466. **31**. Ding XQ et al. J Magn Reson Imaging. 2008;27:998-1004. **32**. Feldmann R et al. Mol Genet Metab. 2019;126:246-249. **33**. Greeves LG et al. Acta Paediatr. 1997;86:242-244. 34. Hawks Z et al. Neuroimage Clin. 2019;23:101916. 35. Hermida-Ameijeiras A et al. Nedicine (Baltimore). 2017;96:e9322. 36. Htun P et al. PLoS One. 2015;10:e0135930. 37. Huemer M et al. Mol Genet Metab. 2008;94:46-51. **38.** Hvas AM et al. *J Inherit Metab Dis.* 2006;29:47-53. **39.** Kono K et al. *Radiology.* 2005;236:630636. **40.** Lage S et al. *J Inherit Metab Dis.* 2010;33 Suppl 3:S363-371. **41.** Laule C et al. *J Magn Reson Imaging.* 2007;26:1117-1121. **42.** Lubout CMA et al. J Inherit Metab Dis. 2020;43:251-258. 43. Millet P et al. Clin Biochem. 2005;38:1127-1132. 44. Modan-Moses D et al. J Inherit Metab Dis. 2007;30:202-208. 45. Nagasaka H et al. J Bone Miner Metab. 2011;29:737-743. 46. Nardecchia F et al. Mol Genet Metab. 2015;115:84-90. 47. Nardecchia F et al. Mol Genet Metab. 2019;128:53-56. 48. Pedersen HE. Acta Neurol Scand. 1974;50:599-610. 49. Peng H et al. J Inherit Metab Dis. 2014;37:237-243. 50. Pérez-Dueñas B et al. Acta Paediatr. 2002;91:899-904 51. Pérez-Dueñas B et al. Neurology. 2006;66:1074-1078. 52. Phillips MD et al. AJNR Am J Neuroradiol. 2001;22:1583-1586. 53. Pietz J et al. Eur J Pediatr. 1998;157:824-830. 54. Pietz J et al. Pediatr Res. 1995;38:657-663. 55. Pietz J et al. J Clin Invest. 1999;103:1169-1178. 56. Porta F et al. J Inherit Metab Dis. 2008;31 Suppl 2:S339-342. 57. Roato I et al. J Hum Nutr Diet. 2013;26 Suppl 1:1-6. 59. Robinson M et al. J Pediatr. 2000;136:545-547. 60. Rocha JC et al. Mol Genet Metab. 2012;107:659-663. 61. Röricht S et al. J Neurol. 1999;246:21-30. 62. Rupp A et al. J Cereb Blood Flow Metab. 2001;21:276-284. 63. Schwahn B et al. Acta Paediatr. 1998;87:61-63. 64. Sirrs SM et al. Radiology. 2007;242:236-243 65. Sumanszki C et al. Ann Nutr Metab. 2019;75:16-23. 66. Vermathen P et al. Magn Reson Med. 2007;58:1145-1156. 67. Vugteveen I et al. Mol Genet Metab. 2011;102:13-17. 68. Weigel C et al. Ann Nutr Metab. 2008;53:91-95. 69. White DA et al. Mol Genet Metab. Metab. 2013;110:213-217. 70. Zeman J et al. J Intellect Disabil Res. 2013;57:567-579. 74. Koch R et al. J Inherit Metab Dis. 2002;25:333-346. 75. Schulz B et al. Acta Paediatr. 1995;84:743-748. 76. Pearsen KD et al. AJNR Am J Neuroradiol. 1991;12:403-406. 78. MacDonald A et al. J Inherit Metab Dis. 2004;27:127-135. 79. Pena MJ et al. Orphanet J Rare Dis. 2021;16:84. 80. Pinto A et al. Eur J Clin Nutr. 2017;71:1230-1234. 81. Das AM et al. J Inherit Metab Dis. 2007;30:209-216. 83. Mirás A et al. Mol Genet Metab

2013;108:149-154. 84. Procházková D et al. Biol Trace Elem Res. 2013;154:178-184. 85. Harding CO et al. Mol Genet Metab. 2018;124:20-26. 86. Lee P et al. Am J Med Genet A. 2008;146a:2851-2859