

Efficacy and safety of valoctocogene roxaparvovec 4 years after gene transfer in GENE8-1


Rauch A¹, Chambost H², Leavitt AD³, Mahlangu J⁴, Raheja P⁵, Symington E⁶, Quon DV⁷, Giermasz A⁸, Kenet G⁹, Lowe G¹⁰, Key NS¹¹, Millar CM^{12,13}, Pipe SW¹⁴, Madan B¹⁵, Chou S-C¹⁶, Klamroth R^{17,18}, Mason J^{19,20}, Peyvandi F^{21,22}, Yu H²³, Robinson TM²³, Ozelo MC²⁴

¹CHU Lille, Hematology Transfusion, Lille, France; ²Department of Pediatric Hematology Oncology, Children Hospital La Timone & Aix Marseille University, Marseille, France; ³Adult Hemophilia Treatment Center, University of California San Francisco, San Francisco, CA, USA; ⁴Hemophilia Comprehensive Care Center, Charlotte Maxeke Johannesburg Academic Hospital, University of the Witwatersrand and NHLS, Johannesburg, South Africa; ⁵The Royal London Hospital Haemophilia Centre, Barts Health NHS Trust, London, UK; ⁶Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK; ⁷Orthopaedic Hemophilia Treatment Center, Los Angeles, CA, USA; ⁸Hemophilia Treatment Center, University of California Davis, Sacramento, CA, USA; ⁹The National Hemophilia Center and Amalia Bron Research Institute of Thrombosis and Hemostasis, Sheba Medical Center, Tel Hashomer, Tel Aviv University, Tel Aviv, Israel; ¹⁰West Midlands Adult Haemophilia Comprehensive Care Centre, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK; ¹¹UNC Blood Research Center, University of North Carolina, Chapel Hill, NC, USA; ¹²Centre for Haematology, Imperial College London, London, UK; ¹³Imperial College Healthcare NHS Trust, London, UK; ¹⁴Departments of Pediatrics and Pathology, University of Michigan, Ann Arbor, MI, USA; ¹⁵Guy's and St Thomas' NHS Foundation Trust, London, UK; ¹⁶Division of Hematology, Department of Internal Medicine, National Taiwan University Hospital, Taipei, Taiwan; ¹⁷Vascular Medicine and Haemostaseology, Vivantes Klinikum im Friedrichshain, Berlin, Germany; ¹⁸Institute of Experimental Hematology and Transfusion Medicine, University Hospital Bonn, Medical Faculty, University of Bonn, Bonn, Germany; ¹⁹Queensland Haemophilia Centre, Cancer Care Services, Royal Brisbane and Women's Hospital, Brisbane, QLD, Australia; ²⁰University of Queensland, Brisbane, QLD, Australia; ²¹Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Angelo Bianchi Bonomi Hemophilia and Thrombosis Center, Milan, Italy; ²²Università degli Studi di Milano, Department of Pathophysiology and Transplantation, Milan, Italy; ²³BioMarin Pharmaceutical Inc., Novato, CA, USA; ²⁴Hemocentro UNICAMP, Department of Internal Medicine, School of Medical Sciences, University of Campinas, Campinas, SP, Brazil

Introduction

Valoctocogene roxaparvovec for severe hemophilia A


- Valoctocogene roxaparvovec (AAV5-hFVIII-SQ) is a liver-directed gene therapy that transfers a B-domain-deleted FVIII coding sequence to enable FVIII production in people with severe hemophilia A (FVIII ≤ 1 IU/dL)^{1,2}
- In the open-label, phase 3 GENE8-1 trial, participants who received 6x10¹³ vg/kg valoctocogene roxaparvovec had improved protection from bleeds compared with regular FVIII prophylaxis over 3 years^{1,2}
- Here, we evaluate efficacy and safety outcomes 4 years after treatment




Methods

Study design

Eligibility	Endpoints
<ul style="list-style-type: none">Adult men with severe hemophilia A (FVIII ≤ 1 IU/dL)Previously receiving FVIII prophylaxisNo history of FVIII inhibitors or anti-AAV5 antibodiesNo significant liver dysfunction, fibrosis, or cirrhosis	<ul style="list-style-type: none">FVIII activityChange from baseline<ul style="list-style-type: none">Annualized bleeding rateAnnualized FVIII infusion rateHRQOL (covered in a separate poster)Safety



Participant disposition



Results

Baseline characteristics

Baseline characteristics	Rollover population N = 112	mITT N = 132	ITT N = 134
Age, years, mean (range)	31.8 (19–70)	31.4 (18–70)	31.7 (18–70)
Race, n (%)			
White	78 (69.6)	94 (71.2)	96 (71.6)
Asian	17 (15.2)	19 (14.4)	19 (14.2)
Black or African American	14 (12.5)	15 (11.4)	15 (11.2)
Hawaiian or Pacific Islander	1 (0.9)	1 (0.8)	1 (0.7)
Not provided	2 (1.8)	3 (2.3)	3 (2.2)
Hispanic or Latino ethnicity, n (%)	5 (4.5)	7 (5.3)	7 (5.2)
BMI, kg/m ² , mean \pm SD	25.2 \pm 4.7	25.3 \pm 4.6	25.3 \pm 4.6
Medical history, n (%)			
Hepatitis B	17 (15.2)	18 (13.6)	20 (14.9)
Hepatitis C	33 (29.5)	39 (29.5)	41 (30.6)
HIV	0	0	2 (1.5)
Number of problem joints,* n (%)			
0	82 (73.2)	95 (72.0)	97 (72.4)
1	13 (11.6)	17 (12.9)	17 (12.7)
2	9 (8.0)	9 (6.8)	9 (6.7)
3	6 (5.4)	8 (6.1)	8 (6.0)
≥ 3	2 (1.8)	3 (2.3)	3 (2.2)

*Problem joints were those with chronic joint pain, chronic synovitis, hemophilic arthropathy, limited motion, or recurrent bleeding. BMI, body mass index; HIV, human immunodeficiency virus; ITT, intention-to-treat; mITT, modified ITT; SD, standard deviation.

Safety (ITT population)

No new safety signals in year 4

- In year 4, no new safety signals were reported
 - Low-grade, transient ALT elevations remained the most common AE in year 4
- No treatment-related SAEs occurred
- No new malignancies were reported
- As of the cutoff date, no FVIII inhibitors were observed and no thromboembolic events occurred

Participants, n (%)		Year 1 (N = 134)	Year 2 (N = 134)	Year 3 (N = 131)	Year 4 (N = 131)	All follow-up
AEs		134 (100.0)	113 (84.3)	105 (80.2)	106 (80.9)	134 (100.0)
SAEs		21 (15.7)	6 (4.5)	9 (6.9)	13 (9.9)	37 (27.6)
Treatment-related AEs*		123 (91.8)	28 (20.9)	15 (11.5)	10 (7.6)	123 (91.8)
Glucocorticoid-related AEs*		80 (59.7)	10 (7.5)	1 (0.8)	1 (0.8)	81 (60.4)
AEs of special interest	ALT elevation	114 (85.1)	40 (29.9)	31 (23.7)	56 (42.7)	121 (90.3)
	ALT elevation grade ≥ 3	11 (8.2)	1 (0.7)	0	1 (0.8) ^b	12 (9.0)
	Potential Hy's law case	0	0	0	0	0
	Infusion-related reactions*	12 (9.0)	0	0	0	12 (9.0)
	Systemic hypersensitivity	7 (5.2)	0	0	0	7 (5.2)
	Anaphylactic or anaphylactoid reactions	3 (2.2)	0	0	0	3 (2.2)
	Thromboembolic events	0	0	0	0	0
Anti-FVIII neutralizing antibodies		0	0	0	0	0
Malignancy (except nonmelanoma skin cancer)		0	0	1 (0.8)	0	1 (0.7)


*Treatment-related and glucocorticoid-related AEs were assessed by the investigator. ^bThis event was downgraded after the data cutoff (November 15, 2023). ^cInfusion-related reactions were defined as AEs occurring during valoctocogene roxaparvovec infusion or within 6 hours post-infusion. AE, adverse event; ALT, alanine aminotransferase; FVIII, factor VIII; SAE, serious AE.

ALT elevation and glucocorticoid use

	During year 4	With AEs in year 4 (N = 131)
In year 4, 56 (42.7%) participants experienced an ALT elevation, most of which were low-grade and transient		
No participants initiated glucocorticoids to manage ALT elevations after week 84		
ALT elevation >ULN, n (%)		21 (16.0)
ALT elevation >1.5x baseline, n (%)		55 (42.0)
Used glucocorticoids for any purpose, n (%)		3 (2.3)
Total duration, weeks, median (range)		1.4 (1.0–12.1)
Total dose, mg, median (range)		200.0 (200–1475)
Used glucocorticoids for ALT elevation, n (%)		0 (0.0)
Total duration, weeks, median (range)		NA
Total dose, mg, median (range)		NA
AE, adverse event; ALT, alanine aminotransferase; NA, not applicable; ULN, upper limit of normal.		


FVIII activity (mITT population)

FVIII activity maintained between years 3 and 4



Because 2 participants did not reach year 4 follow-up, week 208 data are based on 130 participants. For participants who discontinued the study, missing FVIII values post-discontinuation were imputed as 0 IU/dL through the data cutoff date. CSA, chromogenic substrate assay; mITT, modified intention-to-treat; Q, quartile; SE, standard error.

FVIII activity ranges at the end of year 4




Most participants remain in the mild hemophilia range

Because 2 participants did not reach year 4 follow-up, week 208 data are based on 130 participants. For participants who discontinued the study, missing FVIII values post-discontinuation were imputed as 0 IU/dL through the data cutoff date. CSA, chromogenic substrate assay; FVIII, factor VIII; LLOQ, lower limit of quantification.

Annualized bleed rate (Rollover population)

Reduction in treated bleeds maintained over 4 years




ABR for treated bleeds decreased >80% from baseline during the post-prophylaxis period

In year 4, >70% of participants had no treated bleeds

Missing data were not imputed. ABR, annualized bleeding rate; CI, confidence interval; Q, quartile; SD, standard deviation.

Annualized FVIII infusion rate (Rollover population)

Reduction of FVIII infusion rate maintained through year 4




Annualized FVIII infusion rate decreased >95% from baseline during the post-prophylaxis period

Missing data were not imputed. AFR, annualized FVIII infusion rate; CI, confidence interval; Q, quartile; SD, standard deviation.

Return to prophylaxis (ITT population)

Most participants remain off prophylaxis




Overall, 24 participants resumed prophylaxis

Since the previous data cutoff, 7 additional participants resumed prophylaxis

Missing data were not imputed. ABR, annualized bleeding rate; CSA, chromogenic substrate assay; FVIII, factor VIII; RTP, return to prophylaxis.


Conclusions

A single infusion of valoctocogene roxaparvovec provides durable bleeding protection for 4 years with an acceptable safety profile




No new safety signals

- ALT elevation remained the most common AE in year 4; none have required glucocorticoid use since year 2
- No FVIII inhibitors or thromboembolic events



FVIII activity was maintained

- FVIII activity remained in the mild hemophilia range
- Slope of decline in FVIII activity continues to approach 0
- Among the 17 participants dosed ≥ 5 years prior, year 5 values were similar to year 4



Durable hemostatic efficacy

- Rate of treated bleeds in the post-prophylaxis period remains decreased >80% from baseline
- Most participants had no treated bleeds during year 4

Most participants remain off prophylaxis

- Decisions to return to prophylaxis were individual and part of a shared decision-making process that considered multiple factors

References

1. Ozelo M, et al. *N Engl J Med*. 2022;386(11):1013–25. 2. Mahlangu J, et al. *N Engl J Med*. 2023;388:694–705.

Acknowledgements

Thank you to all trial participants, their families, study-site personnel, and investigators. Funding for this study was provided by BioMarin Pharmaceutical Inc. Medical writing support was provided by Amin Ghane, PhD, of AlphaBioCom, a Red Nucleus company, and funded by BioMarin Pharmaceutical Inc. Project management support was provided by Gillian Clague, CMPP, of BioMarin Pharmaceutical Inc.

