

P-583: Efficacy and safety of pegvaliase in adolescents with phenylketonuria: Interim results from PEGASUS part 2

Amarilis Sanchez-Valle¹, Ashley Andrews², Julia B. Hennermann³, Jasmine Knoll⁴, Yuri A. Zarate⁵, Dammie Brown⁶, Konstantia-Maria Chavele⁷, Kristin Lindstrom⁸, Naomi R.M. Schwartz⁶, Janet A. Thomas⁸

¹Morsani College of Medicine, University of South Florida, Tampa, FL, USA; ²University of Utah, Salt Lake City, UT, USA; ³University Medical Center Mainz, Mainz, Germany; ⁴Phoenix Children's Hospital, Phoenix, AZ, USA; ⁵University of Kentucky College of Medicine, Lexington, KY, USA; ⁶BioMarin Pharmaceutical Inc., Novato, CA, USA; ⁷BioMarin (UK) Ltd., London, UK; ⁸University of Colorado School of Medicine, Aurora, CO, USA

Introduction


- Phenylketonuria (PKU) is an autosomal recessive condition characterized by elevated blood phenylalanine (Phe) levels that can lead to neurocognitive, behavioral, and psychiatric symptoms if left unmanaged¹
- Blood Phe levels of ≤360 μmol/L are recommended for all individuals with PKU in the US; European guidelines recommend 120 to 600 μmol/L for adolescents aged 12 to 18 years^{2,3}
- Pegvaliase, a subcutaneous enzyme substitution therapy, lowers blood Phe in adults with PKU^{4,5}
- In PEGASUS (NCT05270837) part 1, the efficacy and safety of pegvaliase in adolescents with PKU (aged 12–17 years) were consistent with outcomes observed in adults, leading to the recent US Food and Drug Administration and European Medicines Agency approvals in this population^{4,6,7}

Methods

Study design

- PEGASUS is a 2-part, randomized (2:1), phase 3 study of efficacy and safety of pegvaliase vs diet-only in adolescents with a confirmed PKU diagnosis (**Figure 1**)
 - After 72 weeks (part 1), participants on diet-only transitioned to pegvaliase for the long-term extension (part 2)

Figure 1. PEGASUS study design



^aParticipants were randomized pegvaliase:diet-only in a 2:1 ratio. EU, European Union; Phe, phenylalanine; PKU, phenylketonuria.

Objective

To evaluate long-term efficacy and safety data from adolescents with PKU randomized to pegvaliase based on an interim analysis of the extension phase (part 2) of PEGASUS


Analysis population

- Only participants initially randomized to pegvaliase in part 1 (N = 36) were included in this analysis

Endpoints

- We report interim part 2 summary efficacy and safety data from an October 2025 data cutoff date (**Figure 2**)

Figure 2. Analysis endpoints



^aAt last assessment before data cutoff (October 2025). AE, adverse event; Phe, phenylalanine; SAE, serious AE.

Results

Participants and treatment exposure

- Baseline mean ± standard deviation (SD) blood Phe was 1025.3 ± 254.1 μmol/L, and mean ± SD age at enrollment was 14.4 ± 1.3 years (**Table 1**)
- At the October 2025 data cutoff, mean ± SD treatment duration was 27.6 ± 8.7 months

Safety

- Hypersensitivity was reported in 34 (94.4%) participants, injection-site reaction in 32 (88.9%), and arthralgia/arthritis in 28 (77.8%; **Table 2**)
- Three (8.3%) participants experienced anaphylaxis (National Institute of Allergy and Infectious Diseases/Food Allergy and Anaphylaxis Network criteria)
- AEs led to treatment and study discontinuation for 2 (5.6%) and 1 (2.8%) participants, respectively

Table 1. Demographics and baseline characteristics

	Pegvaliase N = 36
Age, years, mean ± SD	14.4 ± 1.3
12–15, n (%)	30 (83.3)
16–17, n (%)	6 (16.7)
Sex, n (%)	
Female	22 (61.1)
Male	14 (38.9)
Race, n (%)^a	
White	30 (83.3)
Black or African American	2 (5.6)
Asian	1 (2.8)
Other ^b	0
Blood Phe concentration (μmol/L)	
Mean ± SD	1025.3 ± 254.1
Median (min, max)	996.0 (635.3, 1554.7)
Treatment duration, months	
Mean ± SD	27.6 ± 8.7
Median (Q1, Q3)	28.9 (26.8, 31.8)

^aNot reported, n = 3. ^bOther includes American Indian or Alaska Native, Native Hawaiian or Pacific Islander, and Other. Max, maximum; min, minimum; Phe, phenylalanine; Q, quartile; SD, standard deviation.

Table 2. Incidence of safety events


Participants with AE, n (%)	Pegvaliase N = 36
Any AE	36 (100)
AEs assessed by investigator as related to study drug	36 (100)
AEs leading to dose reduction	11 (30.6)
AEs leading to dose interruption	21 (58.3)
AEs leading to study drug discontinuation	2 (5.6)
AEs leading to study discontinuation	1 (2.8)
Any SAE	5 (13.9)
Most common (≥30%) AEs by preferred term	
Hypersensitivity	34 (94.4)
Injection-site reaction	32 (88.9)
Arthralgia/arthritis	28 (77.8)
Hypersensitivity reaction (US)	18 (50.0)
Injection-site skin reaction ≥14 days duration	17 (47.2)
Hypersensitivity reaction (EU)	16 (44.4)
Arthralgia (steroid treatment)	15 (41.7)
Neuropsychiatric disorder	15 (41.7)
Hypophenylalaninemia on a single measurement	14 (38.9)
Hypophenylalaninemia on 2 or more consecutive measurements	11 (30.6)

AE, adverse event; EU, European Union; SAE, serious AE.

Blood Phe ≤600, ≤360, and ≤120 μmol/L threshold achievement

- Blood Phe ≤600, ≤360, and ≤120 μmol/L was achieved by 25 (69.4%), 20 (55.6%), and 17 (47.2%) participants, respectively
- The median (quartile 1 [Q1], quartile 3 [Q3]) time to ≤600, ≤360, and ≤120 μmol/L threshold achievement was 1.2 (1.0, 11.1), 10.0 (6.7, 14.7), and 13.8 (10.1, 20.1) months, respectively (**Figure 3**)
 - The median (Q1, Q3) dose to reach ≤600 μmol/L blood Phe was 15.0 (2.5, 40.0) mg/day, and the median (Q1, Q3) dose to reach both the ≤360 and ≤120 μmol/L thresholds was 40.0 (20.0, 40.0) mg/day

Figure 3. A) Proportion of participants, B) median time, and C) median dose to achieve blood Phe thresholds



Changes in protein intake from baseline

- For those who achieved blood Phe ≤360 μmol/L while on study, the mean percentage change from baseline in dietary protein intake was –63.4% from medical food and +223.9% from intact food at last assessment (n = 20; **Table 3**)
- For those who achieved blood Phe ≤120 μmol/L while on study, the mean percentage change from baseline in dietary protein intake was –74.1% from medical food and +252.5% from intact food at last assessment (n = 17)
- Notably, as of the data cutoff date, 9 participants discontinued medical food entirely

Table 3. Protein intake at last assessment for participants who achieved blood Phe threshold at any point while on study

	Pegvaliase	
	≤360 μmol/L blood Phe (n = 20)	≤120 μmol/L blood Phe (n = 17)
Medical food		
Mean ± SD, g/kg	0.3 ± 0.4	0.2 ± 0.3
Change from baseline, %	–63.4 ± 46.7	–74.1 ± 41.5
Median (min, max), g/kg	0.2 (0, 1)	0.0 (0, 1)
Change from baseline, %	–82.3 (–100, 59)	–100.0 (–100, 59)
Intact food		
Mean ± SD, g/kg	0.7 ± 0.4	0.8 ± 0.4
Change from baseline, %	223.9 ± 189.9	252.5 ± 191.1
Median (min, max), g/kg	0.6 (0, 2)	0.7 (0, 2)
Change from baseline, %	184.8 (3, 684)	234.3 (3, 684)

Max, maximum; min, minimum; Phe, phenylalanine; SD, standard deviation.

References

1. van Spronsen FJ, et al. *Nat Rev Dis Primers*. 2021;7(1):36. 2. van Wegberg AMJ, et al. *Mol Genet Metab*. 2025;145(2):109125. 3. Smith WE, et al. *Genet Med*. 2025;27(1):101289. 4. PALYNZIQ (pegvaliase-pqpz). Prescribing information. BioMarin Pharmaceutical Inc.; 2026. 5. Thomas J, et al. *Mol Genet Metab*. 2018;124(1):27–38. 6. Sacharow S, et al. Safety and efficacy of pegvaliase in adolescents with phenylketonuria: primary results from PEGASUS, a phase 3 open-label randomized controlled study. Oral presentation at ICiEM 2025; September 2–6, 2025. 7. European Medicines Agency. Palynzic - opinion on variation to marketing authorisation. Accessed June 16, 2026. <https://www.ema.europa.eu/en/medicines/human/variation/palynzic>

Acknowledgments

We thank all trial participants, their families, study-site staff, and investigators. Medical writing support was provided by Rachel R. Corrigan, PhD, of Red Nucleus, and funded by BioMarin Pharmaceutical Inc.

Disclosures

AS-V has received consulting fees from Acer Therapeutics, Applied Therapeutics, BioMarin Pharmaceutical Inc., Horizon, and Ultragenyx and has participated as a clinical trial investigator for Acadia, Alexion, Amicus, BioMarin Pharmaceutical Inc., Cerecor, Genzyme, Homology Medicines, Horizon, Kaleido Biosciences, Mallinckrodt, and Ultragenyx. **AA** has participated in advisory boards for BioMarin Pharmaceutical Inc. **JBH** has received travel support and/or honoraria from Amicus, Chiesi, Immedica, Sanofi, and Takeda. **JK** and **YAZ** have no declarations. **DB, KL,** and **NRMS** are employees and stockholders of BioMarin Pharmaceutical Inc. **K-MC** is an employee and potential stockholder of BioMarin (UK) Ltd. **JAT** has participated as a clinical trial investigator for BioMarin Pharmaceutical Inc., Homology Medicines, Sanofi, and Synlogic and as an advisory board member for BioMarin Pharmaceutical Inc.

To view a copy of this poster, scan this QR code. Copies of this poster obtained through the QR code are for personal use only and may not be reproduced without permission from the authors.

